

SCHWERPUNKT: Entwicklung der Herzmedizin

- **Entwicklungen der Katheterablation von Vorhofflimmern: Von Cox-Maze-Operation zur Pulmonalvenenisolation 2020 (S. 19–26)**

S. Oebel, N. Dagres und G. Hindricks

Literatur:

1. Buist TJ, Zipes DP, Elvan A. Atrial fibrillation ablation strategies and technologies: past, present, and future. *Clin Res Cardiol* 2020 Oct 22. doi: 10.1007/s00392-020-01751-5. Epub ahead of print. PMID: 33089361
2. Calkins H et al. *Europace* 2018 Jan 1; 20(1): e1-e160. doi: 10.1093/europace/eux274. PMID: 29016840; PMCID: PMC5834122
3. Breithardt G et al. Probleme der antiarrhythmischen Therapie bei Vorhofflimmern [Problems with anti-arrhythmia therapy in atrial fibrillation]. *Z Kardiol* 1994; 83 Suppl 5: 63–9. German. PMID: 7846947
4. Dagres N et al. Treating arrhythmias: an experts opinion. *Expert Opin Pharmacother.* 2011 Jun; 12(9): 1359–67. doi: 10.1517/14656566.2011.555397. Epub 2011 Feb 26. PMID: 21352091
5. Camm AJ. Hopes and disappointments with antiarrhythmic drugs. *Int J Cardiol* 2017 Jun 15; 237: 71–74. doi: 10.1016/j.ijcard.2017.03.056. Epub 2017 Mar 16. PMID: 28365182
6. Brugada J et al. 2019 ESC Guidelines for the management of patients with supraventricular tachycardiaThe Task Force for the management of patients with supraventricular tachycardia of the European Society of Cardiology (ESC). *Eur Heart J* 2020 Feb 1; 41(5): 655–720. doi: 10.1093/eurheartj/ehz467. PMID: 31504425
7. Cox JL et al. The surgical treatment of atrial fibrillation. III. Development of a definitive surgical procedure. *J Thorac Cardiovasc Surg* 1991 Apr; 101(4): 569–83. PMID: 2008095
8. Cox JL, Jaquiss RD, Schuessler RB, Boineau JP. Modification of the maze procedure for atrial flutter and atrial fibrillation. II. Surgical technique of the maze III procedure. *J Thorac Cardiovasc Surg* 1995 Aug; 110(2): 485–95. doi: 10.1016/S0022-5223(95)70245-8. PMID: 7637366
9. Cox JL, Schuessler RB, Lappas DG, Boineau JP. An 8 1/2-year clinical experience with surgery for atrial fibrillation. *Ann Surg* 1996 Sep; 224(3): 267–73; discussion 273–5. doi: 10.1097/00000658-199609000-00003. PMID: 8813255; PMCID: PMC1235364
10. Kottkamp H et al. Specific linear left atrial lesions in atrial fibrillation: intraoperative radiofrequency ablation using minimally invasive surgical techniques. *J Am Coll Cardiol* 2002 Aug 7; 40(3): 4–7580. doi: 10.1016/s0735-1097(02)01993-9. PMID: 12142113
11. Tanner H et al. Trigger activity more than three years after left atrial linear ablation without pulmonary vein isolation in patients with atrial fibrillation. *J Am Coll Cardiol* 2005 Jul 19; 46(2): 338–43. doi: 10.1016/j.jacc.2005.03.063. PMID: 16022965
12. Doll N et al. Die chirurgische Behandlung des therapierefraktären Vorhofflimmerns [Surgical treatment of therapy resistant atrial fibrillation]. *Z Kardiol* 2003 Sep; 92(9): 712–20. German. doi: 10.1007/s00392-003-0953-0. PMID: 14508587
13. Swartz JF et al. A catheter-based curative approach to atrial fibrillation in humans, *Circulation* 1994; vol. 90 (pg. I-335) (Abstract)
14. Haissaguerre M et al. Spontaneous initiation of atrial fibrillation by ectopic beats originating in the pulmonary veins. *N Engl J Med* 1998 Sep 3; 339(10): 659–66. doi: 10.1056/NEJM199809033391003. PMID: 9725923

15. Haïssaguerre M et al. Electrophysiological end point for catheter ablation of atrial fibrillation initiated from multiple pulmonary venous foci. *Circulation* 2000 Mar 28; 101(12): 1409–17. doi: 10.1161/01.cir.101.12.1409. PMID: 10736285
16. Oral H et al. Circumferential pulmonary-vein ablation for chronic atrial fibrillation. *N Engl J Med* 2006 Mar 2; 354(9): 934–41. doi: 10.1056/NEJMoa050955. PMID: 16510747
17. Ouyang F et al. Electrophysiological findings during ablation of persistent atrial fibrillation with electroanatomic mapping and double Lasso catheter technique. *Circulation* 2005 Nov 15; 112(20): 3038–48. doi: 10.1161/CIRCULATIONAHA.105.561183. Epub 2005 Nov 7. PMID: 16275866
18. Clarnette JA et al. Outcomes of persistent and long-standing persistent atrial fibrillation ablation: a systematic review and meta-analysis. *Europace* 2018 Nov 1; 20(FI_3): f366-f376. doi: 10.1093/europace/eux297. PMID: 29267853
19. Kircher S et al. Individually tailored vs. standardized substrate modification during radiofrequency catheter ablation for atrial fibrillation: a randomized study. *Europace* 2018 Nov 1; 20(11): 1766–1775. doi: 10.1093/europace/eux310. PMID: 29177475
20. Kuck KH et al. Cryoballoon or Radiofrequency Ablation for Paroxysmal Atrial Fibrillation. *N Engl J Med* 2016 Jun 9; 374(23): 2235–45. doi: 10.1056/NEJMoa1602014. Epub 2016 Apr 4. PMID: 27042964
21. Bollmann A et al. Cardiac tamponade in catheter ablation of atrial fibrillation: German-wide analysis of 21 141 procedures in the Helios atrial fibrillation ablation registry (SAFER). *Europace* 2018 Dec 1; 20(12): 1944–1951. doi: 10.1093/europace/euy131. PMID: 29982554
22. Arbelo E et al. Contemporary management of patients undergoing atrial fibrillation ablation: in-hospital and 1-year follow-up findings from the ESC-EHRA atrial fibrillation ablation long-term registry. *Eur Heart J* 2017 May 1; 38(17): 1303–1316. doi: 10.1093/eurheartj/ehw564. PMID: 28104790
23. Dagres N et al. Complications of atrial fibrillation ablation in a high-volume center in 1,000 procedures: still cause for concern? *J Cardiovasc Electrophysiol* 2009 Sep; 20(9): 1014–9. doi: 10.1111/j.1540-8167.2009.01493.x. Epub 2009 May 20. PMID: 19490383
24. Schoene K et al. Acquired Pulmonary Vein Stenosis After Radiofrequency Ablation for Atrial Fibrillation: Single-Center Experience in Catheter Interventional Treatment. *JACC Cardiovasc Interv* 2018 Aug 27; 11(16): 1626–1632. doi: 10.1016/j.jcin.2018.05.016. PMID: 30139470
25. Halm U et al. Thermal esophageal lesions after radiofrequency catheter ablation of left atrial arrhythmias. *Am J Gastroenterol* 2010 Mar; 105(3): 551–6. doi: 10.1038/ajg.2009.625. Epub 2009 Nov 3. PMID: 19888201
26. Hindricks G et al. 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association of Cardio-Thoracic Surgery (EACTS). *Eur J Heart* 2020 Aug 29: ehaa612. doi: 10.1093/eurheartj/ehaa612. Epub ahead of print. PMID: 32860505
27. Kotadia ID, Williams SE, O'Neill M. High-power, Short-duration Radiofrequency Ablation for the Treatment of AF. *Arrhythm Electrophysiol Rev* 2020; 8(4): 265–272. doi:10.15420/aer.2019.09
28. Sugrue A et al. Irreversible electroporation for the treatment of cardiac arrhythmias. *Expert Rev Cardiovasc Ther* 2018 May; 16(5): 349–360. doi: 10.1080/14779072.2018.1459185. PMID: 29595355
29. Piccini JP et al. Changes in quality of life, cognition and functional status following catheter ablation of atrial fibrillation. *Heart* 2020 Oct 12: heartjnl-2020-316612. doi: 10.1136/heartjnl-2020-316612. Epub ahead of print. PMID: 33046527
30. Raatikainen MJ et al. Radiofrequency catheter ablation maintains its efficacy better than antiarrhythmic medication in patients with paroxysmal atrial fibrillation: On-treatment analysis

- of the randomized controlled MANTRA-PAF trial. *Int J Cardiol* 2015 Nov 1; 198: 108–14. doi: 10.1016/j.ijcard.2015.06.160. Epub 2015 Jul 4. PMID: 26163901
31. Packer DL. Effect of Catheter Ablation vs Antiarrhythmic Drug Therapy on Mortality, Stroke, Bleeding, and Cardiac Arrest Among Patients With Atrial Fibrillation: The CABANA Randomized Clinical Trial. *JAMA* 2019 Apr 2; 321(13): 1261–1274. doi: 10.1001/jama.2019.0693. PMID: 30874766; PMCID: PMC6450284
32. Richter S, Di Biase L, Hindricks G. Atrial fibrillation ablation in heart failure. *Eur Heart J* 2019 Feb 21; 40(8): 663–671. doi: 10.1093/eurheartj/ehy778. PMID: 30561633
33. Kirchhof P et al. Early Rhythm-Control Therapy in Patients with Atrial Fibrillation. *N Engl J Med* 2020 Oct 1; 383(14): 1305–1316. doi: 10.1056/NEJMoa2019422. Epub 2020 Aug 29. PMID: 32865375
34. Pranata R et al. Longer diagnosis-to-ablation time is associated with recurrence of atrial fibrillation after catheter ablation—Systematic review and meta-analysis. *J Arrhythm* 2019 Dec 27; 36(2): 289–294. doi: 10.1002/joa3. 12294. PMID: 32256876; PMCID: PMC7132183
35. Eitel C et al. Successful nonsurgical treatment of esophagopericardial fistulas after atrial fibrillation catheter ablation: a case series. *Circ Arrhythm Electrophysiol* 2013 Aug; 6(4): 675–81
36. Schoene K et al. Complex cases of acquired pulmonary vein stenosis after radiofrequency ablation: is surgical repair an option? *Europace* 2019; 21(1): 73–79

- **Teil II: Sport – „Polypill“ gegen kardiovaskuläre Risikofaktoren (S. 51–58)**

R. G. Ketelhut, K. Ketelhut

Literatur:

1. Pedisic Z et al. Is running associated with a lower risk of all-cause, cardiovascular and cancer mortality, and is the more the better? A systematic review and meta-analysis. *Br J Sports Med* 2019. <http://dx.doi.org/10.1136/bjsports-2018-100493>
2. Liu X et al. Dose-Response Association between Physical Activity and Incident Hypertension: A Systematic Review and Meta-Analysis of Cohort Studies. *Hypertension* 2017; 69: 813–820
3. Ketelhut K et al. Two years of regular exercise decreases blood pressure and improves motor skills in early childhood. *Sport Sci Health* 2018; DOI 10.1007/s11332-018-0463-0
4. Ketelhut RG et al. Regular exercise as an effective longterm approach in antihypertensive therapy. *Med Sci Sports Exerc* 2004; 1: 4–8
5. Naci H et al. How does exercise treatment compare with antihypertensive medications? A network meta-analysis of 391 randomised controlled trials assessing exercise and medication effects on systolic blood pressure. *Br J Sports Med* 2018. <http://dx.doi.org/10.1136/bjsports-2018-099921>
6. Ketelhut S, Ketelhut RG. The Physiological Responses to Exercise - Type of exercise training and training methods. In: Xiao J (ed.) *Physical Exercise for Human Health*. Springer Nature 2020 (in Druck)
7. Ketelhut RG. Sport bei Hypertonie. *Diabetol* 2015; 11: 394–399
8. Cornelissen VA, Smart NA. Exercise training for blood pressure: a systematic review and meta-analysis. *Am Heart Assoc* 2013; 2: e004473
9. The Task force for the management of arterial hypertension of the European Society of Cardiology and the European Society of Hypertension. 2018 ESC/ESH Guidelines for the management of arterial hypertension. *J Hypertens* 2018; 36: 1953–2041

10. Cornelissen VA, Fagard RH. Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors. *J Hypertens* 2005; 46: 667–675
11. Cornelissen VA, Fagard RH. Effect of resistance training on resting blood pressure: a meta-analysis of randomized controlled trials. *J Hypertens* 2005; 23: 251–259
12. Inder JD et al. Isometric exercise training for blood pressure management: a systematic review and meta-analysis to optimize benefit. *Hypertens Res* 2016; 39: 88–94
13. Mora-Rodriguez R et al. Time-course effects of aerobic interval training and detraining in patients with metabolic syndrome. *Nutr Metab Cardiovasc Dis* 2014; 24(7): 792–798
14. Gillen JB et al. Three minutes of all-out intermittent exercise per week increases skeletal muscle oxidative capacity and improves cardiometabolic health. *PLoS One* 2014; 9(11): e111489
15. Morales-Palomo F et al. Effects of repeated yearly exposure to exercise-training on blood pressure and metabolic syndrome evolution. *J Hypertens* 2017; 35(10): 1992–1999
16. Soltani M et al. High-intensity interval training irrespective of its intensity improves markers of blood fluidity in hypertensive patients. *Clin Exp Hypertens.* 2019 Jul 30: 1–6
17. Martinez Aguirre-Betolaza A. et al. Effects of different aerobic exercise programs on cardiac autonomic modulation and hemodynamics in hypertension: data from EXERDIET-HTA randomized trial. *J Hum Hypertens* 2020; doi:10.1038/s41371-020-0298-4
18. Ciolac EG et al. Effects of high-intensity aerobic interval training vs. moderate exercise on hemodynamic, metabolic and neuro-humoral abnormalities of young normotensive women at high familial risk for hypertension. *Hypertens Res* 2010; 33: 836–843
19. Tjønna AE et al. Aerobic interval training versus continuous moderate exercise as a treatment for the metabolic syndrome. A pilot study. *Circulation* 2008; 118: 346–354
20. Ketelhut S. A comparison of exergaming and aerobic training on systolic and diastolic arterial pressure. (Publikation in Vorbereitung)
21. Ketelhut RG, Ketelhut K. Sport/körperliche Aktivität und Blutdruck. *Herzmed* 2018; 5: 30–38
22. Ketelhut RG et al. Fitness in the fit: does physical conditioning affect cardiovascular risk factors in middle-aged marathon runners? *Eur Heart J* 1996; 17: 199–203
23. Dufaux B et al. Plasmalipoproteins and physical activity: a review. *Int J Sports Med* 1982; 3: 123
24. Wang Y, Xu D. Effects of aerobic exercise on lipids and lipoproteins. *Lipids Health Dis* 2017; 16: 132
25. Christensen R et. al. Effect of Aerobic and Resistance Exercise on Cardiac Adipose Tissues - Secondary Analyses from a Randomized Clinical Trial. *JAMA Cardiol* 2019; 4(8): 778–787
26. Alvarez C et al. High-Intensity Interval Training as a Tool for Counteracting Dyslipidemia in Women. *Int J Sports Med* 2018; 39(05): 397–406
27. Shaw JE et al. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res Clin Pract* 2010; 87: 4–14
28. Colberg SR et al. Exercise and type 2 diabetes: American College of Sports Medicine and the American Diabetes Association: joint position statement. Exercise and type 2 diabetes. *Med Sci Sports Exerc* 2010; 42: 2282–2303
29. Dunstan DW et al. Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diab Care* 2012; 35: 976–983
30. Bailey DP, Locke CD. Breaking up prolonged sitting with light-intensity walking improves postprandial glycemia but breaking up sitting with standing does not. *J Sci Med Sport* 2015; 18: 294–298
31. Maylor BD et al. Effects of Frequency and Duration of Interrupting Sitting on Cardiometabolic Risk Markers. *Int J Sports Med* 2019; 40: 818–824
32. Eriksen L et al. Comparison of the effect of multiple short-duration with single long-duration exercise sessions on glucose homeostasis in type 2 diabetes mellitus. *Diab News* 2008; 1: 5

33. Di Loreto C et al. Make Your Diabetic Patients Walk - Long-term impact of different amounts of physical activity on type 2 diabetes. *Diabetes Care* 2005; 28: 1295–1302
34. Church TS et al. Effects of aerobic and resistance training on hemoglobin A1c levels in patients with type 2 diabetes: a randomized controlled trial. *JAMA* 2010; 304(20): 2253–2262
35. Karstoft K et al. The effects of free-living interval-walking training on glycemic control, body composition, and physical fitness in type 2 diabetic patients: a randomized, controlled trial. *Diabetes Care* 2013; 36(2): 228–36
36. Metcalfe RS et al. Towards the minimal amount of exercise for improving metabolic health: beneficial effects of reduced-exertion high-intensity interval training. *Eur J Appl Physiol* 2012; 112: 2767–2775
37. Little JP et al. Low-volume high-intensity interval training reduces hyperglycemia and increases muscle mitochondrial capacity in patients with type 2 diabetes. *J Appl Physiol* 1985; 2011; 111: 1554–1560
38. Karstoft K et al. The effects of free-living interval-walking training on glycemic control, body composition, and physical fitness in type 2 diabetic patients: a randomized, controlled trial. *Diabetes Care* 2013; 36(2): 228–236
39. Alvarez C et al. Low-Volume High-Intensity Interval Training as a Therapy for Type 2 Diabetes *Int J Sports Med* 2016; 37: 723–729
40. Saris WH et al. How much physical activity is enough to prevent unhealthy weight gain? Outcome of the IASO 1st Stock Conference and consensus statement. *Obes Res* 2003; 4: 101–114
41. Ketelhut K et al. Two years of regular exercise decreases blood pressure and improves motor skills in early childhood. *Sport Sci Health* 2018; DOI 10.1007/s11332-018-0463-0
42. Martin CK et al. Effect of different doses of supervised exercise on food intake, metabolism, and non-exercise physical activity: The E-MECHANIC randomized controlled trial. *Am J Clin Nutr* 2019; 110(3): 583–592
43. Ross R et al. Effects of Exercise Amount and Intensity on Abdominal Obesity and Glucose Tolerance in Obese Adults: A Randomized Trial. *Ann Intern Med* 2015; 162(5): 325–34
44. Graf C, Dordel S. Das CHILT-I-Projekt (Children's Health Interventional Trial). *Bundesgesundheitsblatt* 2011; 54: 313–321
45. Ferguson EW et al. Effects of exercise and conditioning on clotting and fibrinolytic activity in men. *J Appl Physiol* 1987; 62: 1416–1421
46. Szymanski LA et al. Effects of maximal exercise and venous occlusion on fibrinolytic activity in physically active and inactive men. *J Appl Physiol* 1994; 77: 2305–2319
47. Ketelhut RG et al. White blood cell and platelet count: relation to arterial pressure and fitness in normotensive subjects. *DMW* 1999; 3: 90
48. Warlow CP et al. Effect of Exercise on Platelet Count, Adhesion, and Aggregation. *Acta Haematol* 1974; 52: 47–52
49. Ketelhut RG et al. Regelmäßiges Ausdauertraining senkt den Blutdruck und das Fibrinogen bei untrainierten Hypertonikern. *Z Sportmed*, 2003; 7/8: S92
50. Kupchak BR et al. Beneficial effects of habitual resistance exercise training on coagulation and fibrinolytic responses. *Thromb Res* 2013; Jun; 131(6): e227–34
51. da Cunha Nascimento D et al. The interactions between hemostasis and resistance training: a review. *Int J Gen Med* 2012; 5: 249–254
52. Ahmadizad S et al. Effects of time of day and acute resistance exercise on platelet activation and function. *Clin Hemorheol Microcirc* 2010; 45(2–4): 391–399
53. Karampour S, Gaeini AA. Response of coagulation and anti-coagulant factors of elite athletes following acute resistance and high-intensity interval training. *J Sports Med Phys Fitness* 2018 Jan-Feb; 58(1-2): 120–126

54. Valizadeh R et al. The Effect of High Intensity Interval Training on the Response of Coagulation and Fibrinolytic Factors of Hypertensive Patients to One Bout Submaximal Endurance Exercise Report of Health Care 2017; 3(3): 46–57
55. Fiuzo-Luces C et al. Exercise benefits in cardiovascular disease: beyond attenuation of traditional risk factors. Nat Rev Cardiol 2018; 15: 731–43
56. Wen CP et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. Lancet 2011; 378: 1244–53
57. Moholdt T et al. Patterns of physical activity over 22 years and mortality: the HUNT Study, Norway. ESC Congress 2019; 627